

Review of Interactions Between Host Plants, Aphids, Primary Parasitoids and Hyperparasitoids in Vegetable and Cereal Ecosystems in Slovenia

Katarina KOS¹ Olivera PETROVIĆ-OBRADOVIĆ² Vladimir ŽIKIĆ³
Andjeljko PETROVIĆ⁴ Stanislav TRDAN¹ Željko TOMANOVIĆ⁴

¹Department of Agronomy, Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, SI-1111 Ljubljana, SLOVENIA; e-mails: katarina.kos@bf.uni-lj.si, stanislav.trdan@bf.uni-lj.si

²Department of Plant Protection, Faculty of Agriculture, University of Belgrade, Nemanjina 6, SER-11081 Zemun, SERBIA; e-mail: petrovic@agrif.bg.ac.rs

³Department of Biology and Ecology, Faculty of Sciences and Mathematics, University of Niš, Višegradska 33, SER-18000 Niš, SERBIA; e-mail: vzikic@yahoo.com

⁴Institute of Zoology, Faculty of Biology, University of Belgrade, Studentski trg 16, SER-11000 Belgrade, SERBIA; e-mails: andjeljko@bio.bg.ac.rs, ztoman@bio.bg.ac.rs

ABSTRACT

Sampling of aphids and parasitoids was conducted in diverse cereal and vegetable crop habitats in Slovenia in the period from 2006 to 2010. Over 330 samples containing primary or secondary parasitoids or both were taken. In 25 different vegetable crops 15 species of aphids parasitized by 18 species of primary parasitoids were found, and were associated with 13 species of secondary or hyperparasitoids. Eight species of primary parasitoids emerged from mummies of three species of cereal aphids. *Aphidius ervi* Haliday, *A. rhopalosiphi* de Stefani-Perez, *A. uzbekistanicus* Luzhetski, and *Ephedrus plagiator* (Nees), are generally present in cereal crops and most abundant primary parasitoids in cereal crops in Slovenia.

Nine species of hyperparasitoids were found both in cereal and vegetable crops, while four species, i.e. *Alloxysta fulviceps* Curtis, *Coruna clavata* Walker, *Dendrocerus laticeps* Hedicke and *Pachyneuron formosum* Walker, were found only in vegetable crops and two, *Alloxysta brachyptera* Hartig and *Phaenoglyphis villosa* Hartig, only in cereal. 30% of all hyperparasitoid species belonged to genus *Alloxysta*. Cereal and vegetable crops reveal a relatively diverse aphid parasitoid fauna that corresponds to diverse habitat, landscape and vegetation complexity in Slovenia.

Key words: Aphid parasitoid, interactions, vegetable ecosystem, cereal ecosystem, Slovenia.

INTRODUCTION

Insect parasitoids are one of the most diverse and abundant group of arthropods in terrestrial ecosystems (Boivin and Brodeur, 2006) and are widely used as agents in biological control of many agricultural and forest insect pests all over the world. Conservation and augmentation biological control methods require not only to introduce natural enemy, but also to assure suitable environment to enhance the

survival, fecundity, longevity, and behavior of natural enemies to increase their effectiveness. It can also provide them additional or alternative food and hosts/prey, shelter, and protection from adverse conditions (Landis *et al.*, 2000; van den Bosch and Telford, 1964).

Aphids represent very important group of plant pests with high reproductive potential through parthenogenesis and viviparity (Agarwala and Das, 2012). They are causing direct (sucking) and indirect (transmission of viruses and honeydew secretion) damage (Minks and Harrewijn, 1988) on cultivated and wild-growing plants. The populations of aphids can be controlled in different ways, but because of the great damage they cause the producers of plant food, ornamental plants and feed control them mostly with synthetic insecticides (Minks and Harrewijn, 1988; Parker *et al.*, 2006). Over the last decades the release of insect parasitoids in biological control programs has increased all over the world against phytophagous pests in open fields, greenhouses and even in forestry (Boller *et al.*, 2006; Wajnberg, 2010), i.e. cereal aphid parasitoids in Latin America (Zuñiga, 1986), *Aphidius ervi* Haliday to control *Aulacorthum solani* (Kaltenbach) in greenhouses (Henry *et al.*, 2010), and introduction of *TTorymus sinensis* Kamijo in Italy to control chestnut gall wasp *Dryocosmus kuriphilus* Yasumatsu (Quacchia *et al.*, 2008).

The braconid aphid parasitoids are usually oligophagous or polyphagous, attacking more than one host species (Starý, 1970, 1988). However, female parasitoids show some host preferences based on host recognition cues and chemical volatile cues from host plants (Dicke and Sabelis, 1988; Micha *et al.*, 2000; Stoerck *et al.*, 2000; Turlings *et al.*, 1990). The majority of adult parasitoids require some polysaccharide source, like floral nectar or honeydew, to cover their energetic needs required to find hosts (Bianchi and Wäckers, 2008; Godfray, 1994). In addition, some parasitoids, for example synovigenic species, feed on their hosts to get nutrients needed to mature eggs (Godfray, 1994).

In nature, trophic relationships among organisms comprise an extensive web of interactions extending across several trophic levels. The host plants are active trophic level in tritrophic and multitrophic systems directly affecting the ecology and behavior of herbivores and their natural enemies (Sullivan and Völkl, 1999; Tscharntke and Hawkins, 2002). Herbivore feeding damage stimulates host plants to release semiochemicals that can act as attractant or repellent/deterrent to higher trophic levels and thus serve as a plant defense mechanism (Dicke and Sabelis, 1988; Price, 1984).

Insect hyperparasitism can be defined as a highly evolved fourth trophic level relationship that exists between entomophagous insects. As in primary parasitism endo- and ectoparasitism, and idio- and koinobiont life-history traits are known also in hyperparasitism (Strand, 2000). Hyperparasitoids have huge impact on the dynamics of arthropod communities, but the outcome of their introduction in biological control programs is still uncertain. If a facultative hyperparasitoid would have a preference for herbivores rather than for primary parasitoids, biological control could be enhanced (Boivin and Brodeur, 2006). Aphid endohyperparasitoids can use volatile compounds

emitted by the aphid attacked plant for host finding (similar as primary parasitoids), whereas aphid ectohyperparasitoids search randomly for host and do not use specific cues (Blande *et al.*, 2007; Sullivan and Völkl, 1999).

Although aphids represent one of the main pests in arable crops in Slovenia (Modic and Urek, 2008), there was little information about parasitoid species present (Kos *et al.*, 2008, 2009; Milevoj, 2001). Milevoj (2001) mentioned two species, *Aphidius matricariae* Haliday and *Diaeretiella rapae* M'Intosh as native aphid parasitoids found in spring barley. The survey of the tritrophic associations of aphid parasitoids from wide range of habitats in Slovenia was presented by Kos *et al.* in 2012 and can serve as a model to the use of four-trophic studies. Data about hyperparasitoids are generally scarce, and are completely lacking from Slovenia and surrounding countries. In last five years the faunistic composition and abundance of four-trophic relations of aphids, aphidiine parasitoids and hyperparasitoids in vegetable and cereal agroecosystems and nearby non-crop ecosystems was examined in Slovenia. In present study plant-aphid-primary parasitoid-hyperparasitoid interactions on crop plants in vegetable and cereal agroecosystems in Slovenia were determined. The aim of this work is to realize the patterns of host associations (host plant-aphid-primary parasitoid-hyperparasitoid) and to explore the possibilities of aphid parasitoid introduction in biological control programs.

MATERIAL AND METHODS

Sampling of aphids and parasitoids was conducted in diverse cereal and vegetable crop habitats in Slovenia in the period from 2006 to 2010. The collected samples of aphids in crops originated from all over Slovenia, which has three major types of climate: Alpine, Continental and Mediterranean, and is divided on four macro-regions: Alpine, Pannonian, Dinaric, and Mediterranean macro-region. Over 330 samples containing primary or secondary parasitoids or both were taken. For this research only samples of crop plants in vegetable and cereal/corn ecosystems were chosen.

Samples of plants infested with aphid colonies containing also mummified aphids were collected in the field and transferred into transparent plastic containers covered with nylon mesh. Vouchers of aphid adults from each sample were separated, preserved in 70% ethanol, and identified. When necessary, plants were also preserved as herbarium specimens for identification. The remaining aphid samples were maintained in air-conditioned rooms and checked daily for emerging parasitoid adults (Kavallieratos *et al.*, 2004; 2005). The primary parasitoids and hyperparasitoids were then stored in 70% ethanol and later identified at the species level using Olympus SZX 9 stereomicroscope (Olympus Optical Co., Japan) and several published identification keys (Kavallieratos *et al.*, 2001; 2005). Identification of aphids was performed on the Faculty of Agriculture in Zemun (Serbia), identification of parasitoids on the Faculty of Biology in Belgrade (Serbia), and the identification of host plants on Biotechnical Faculty in Ljubljana (Slovenia).

RESULTS

Fifteen species of aphids were found in samples taken from 25 different vegetable crops (belonging to 9 families). They were parasitized by 18 species of primary parasitoids and associated with 13 species of secondary/hyperparasitoids (Table 1). The most polyphagous species was *Lysiphlebus fabarum* (Marshall) with 5 aphid hosts in 8 vegetables from 5 different families (Asteraceae, Chenopodiaceae, Cucurbitaceae, Fabaceae, and Solanaceae), followed by *Aphidius matricariae*, associated with 4 aphid hosts in 8 crop plants from 5 families (Apiaceae, Asteraceae, Brassicaceae, Solanaceae, and Valerianaceae). We have recorded 54 vegetable plant-host aphid-primary parasitoid tritrophic associations.

Also four aphelinid primary parasitoid species were detected in vegetable crops, *Aphelinus chaonia* Walker parasitizing *Aphis fabae* Scopoli in bean and *Uroleucon cichorii* (Koch) in chicory, where *Aphelinus daucicola* Kurdjumov was also found, *Aphelinus mali* Haldeman on *Myzus persicae* (Sulzer) on pepper, and *Aphelinus varipes* Foerster on *U. cichorii* and *Aphis intybi* (Koch) in chicory.

Four different cereals (*Avena sativa* L., *Hordeum vulgare* L., *Triticum aestivum* L. emend. Fiori et Paol., and *Zea mays* L.) most commonly grown in Slovenia were observed for the investigation of cereal aphid and aphid parasitoid/hyperparasitoid fauna. Eight species of primary parasitoids emerged from mummies of three aphid species. *Sitobion avenae* (F.) was the most common, while *Rhopalosiphum maidis* (Fitch), and *R. padi* (L.) were sampled only in maize and barley, respectively (Table 2). Four species of primary parasitoids, *Aphidius ervi*, *A. rhopalosiphii* de Stefani-Perez, *A. uzbekistanicus* Luzhetski, and *Ephedrus plagiator* (Nees) are generally present and most abundant species in cereal crops in Slovenia (Fig. 1). The most frequently sampled hyperparasitoid species were *Alloxysta brevis* Thomson, *A. victrix* Westwood, *Dendrocerus carpenteri* Curtis and *Phaenoglyphis villosa* Hartig. Relations on trophic levels in cereal crops in Slovenia are presented in Figure 1.

Three aphelinid primary parasitoid species were also found in cereal crops, *Aphelinus asychis* Walker parasitizing *S. avenae* in barley, *A. chaonia* parasitizing *S. avenae* in oat and *A. varipes* parasitizing *R. padi* in maize.

Fifteen species of hyperparasitoids from 7 different genera were found in our study (Table 3). Four species were found only in vegetable crops, i.e. *Alloxysta fulviceps* Curtis, *Coruna clavata* Walker, *Dendrocerus laticeps* Hedicke and *Pachyneuron formosum* Walker and two, *Alloxysta brachyptera* Hartig and *Phaenoglyphis villosa*, only in cereal. 30% of all hyperparasitoid species belonged to *Alloxysta* genus. *Alloxysta victrix* found in both crop groups was the most common. 26 specimens of hyperparasitoid *Syrphophagus aphidivorus* Mayr were found on vegetable plants and only two in cereals, while *Asaphes suspensus* Nees, *Dendrocerus carpenteri* and *Pachyneuron aphidis* Bouché species were more numerous in cereal than in vegetable crops.

Review of Interactions Between Host Plants, Aphids, Primary Parasitoids

Table 1. Review of trophic associations (host plant – host aphid – primary parasitoid – secondary parasitoid/hyperparasitoid) on some vegetable crop plants in Slovenia.

Host plant	Host aphid	Primary parasitoid	Secondary parasitoid
<i>Allium cepa</i> L.	<i>Aphis</i> sp.	<i>Aphidius ervi</i> Haliday	<i>Alloxysta victrix</i> Westwood
	<i>Acyrtosiphon pisum</i> (Harris)	<i>Praon</i> sp.	<i>Pachyneuron aphidis</i> Bouché
	<i>Hyperomyzus lactucae</i> (L.)		
<i>Allium porrum</i> L.	<i>Aphis</i> sp.	<i>Aphidius ervi</i> Haliday	<i>Asaphes suspensus</i> Nees
			<i>Dendrocerus carpenteri</i> Curtis
			<i>Dendrocerus laticeps</i> Hedicke
			<i>Pachyneuron muscarum</i> L.
<i>Apium graveolens</i> L.	<i>Cavariella aegopodii</i> Scopoli	<i>Aphidius salicis</i> Haliday	
<i>Beta vulgaris</i> ssp. <i>cicla</i> L.	<i>Aphis fabae</i> Scopoli	<i>Lysiphlebus fabarum</i> (Marshall)	<i>Syrphophagus aphidivorus</i> Mayr
<i>Brassica oleracea</i> L. var. <i>botrytis</i> L. <i>cauliflora</i>	<i>Brevicoryne brassicae</i> (L.)	<i>Diaeretiella rapae</i> M'Intosh	<i>Alloxysta victrix</i> Westwood
<i>Brassica oleracea</i> L. var. <i>capitata</i> L.	<i>Brevicoryne brassicae</i> (L.)	<i>Diaeretiella rapae</i> M'Intosh	<i>Pachyneuron aphidis</i> Bouché
	<i>Lipaphis erysimi</i> (Kaltenbach)	<i>Aphidius matricariae</i> Haliday	<i>Alloxysta victrix</i> Westwood
		<i>Praon volucre</i> (Haliday)	<i>Coruna clavata</i> Walker
			<i>Alloxysta brevis</i> Thomson
			<i>Alloxysta fulviceps</i> Curtis
<i>Brassica oleracea</i> var. <i>gemmifera</i> DC.	<i>Brevicoryne brassicae</i> (L.)	<i>Diaeretiella rapae</i> M'Intosh	
<i>Brassica oleracea</i> L. var. <i>gongyloides</i> L.	<i>Brevicoryne brassicae</i> (L.)	<i>Praon volucre</i> (Haliday)	
<i>Brassica oleracea</i> L. var. <i>italica</i> Plenck	<i>Brevicoryne brassicae</i> (L.)	<i>Diaeretiella rapae</i> M'Intosh	
<i>Brassica pekinensis</i> (Lour.) Rupr.	<i>Myzus persicae</i> (Sulzer)	<i>Aphidius matricariae</i> Haliday	
<i>Capsicum annuum</i> L.	<i>Myzus persicae</i> (Sulzer)	<i>Aphidius matricariae</i> Haliday	
	<i>Aphis nasturtii</i> (Kaltenbach)	<i>Lysiphlebus fabarum</i> (Marshall)	
	<i>Aphis</i> sp.		
	<i>Aulacorthum solani</i> (Kaltenbach)		
<i>Carum carvi</i> L.	<i>Cavariella aegopodii</i> Scopoli	<i>Aphidius salicis</i> Haliday	
		<i>Binodoxys heraclei</i> (Haliday)	
<i>Cichorium intybus</i> L. var. <i>foliosum</i> Hegi	<i>Uroleucon cichorii</i> (Koch)	<i>Aphidius funebris</i> Mackauer	<i>Alloxysta brevis</i> Thomson
	<i>Aphis intybi</i> (Koch)	<i>Aphidius matricariae</i> Haliday	<i>Syrphophagus aphidivorus</i> Mayr
	<i>Myzus persicae</i> (Sulzer)	<i>Lipolexis gracilis</i> (Forster)	
		<i>Lysiphlebus fabarum</i> (Marshall)	
		<i>Praon volucre</i> (Haliday)	
		<i>Praon yomenae</i> Takada	
<i>Cucumis melo</i> L.	<i>Aphis gossypii</i> Glover	<i>Lysiphlebus fabarum</i> (Marshall)	<i>Dendrocerus carpenteri</i> Curtis

Table 1. (Continued)

Host plant	Host aphid	Primary parasitoid	Secondary parasitoid
<i>Cucumis sativus</i> L.	<i>Aphis gossypii</i> Glover	<i>Lysiphlebus fabarum</i> (Marshall)	<i>Alloxysta brevis</i> Thomson
		<i>Binodoxys acalephae</i> (Marshall)	
		<i>Binodoxys angelicae</i> (Haliday)	
<i>Cucurbita pepo</i> L.	<i>Aphis nasturtii</i> (Kaltenbach)	<i>Binodoxys angelicae</i> (Haliday)	<i>Syrphophagus aphidivorus</i> Mayr
<i>Cynara scolymus</i> L.	<i>Brachycaudus cardui</i> (L.)	<i>Diaeretiella rapae</i> M'Intosh	
	<i>Aphis</i> sp.	<i>Lysiphlebus fabarum</i> (Marshall)	
	<i>Pleotrichophorus</i> sp.		
<i>Daucus carota</i> L.	<i>Cavariella aegopodii</i> Scopoli	<i>Ephedrus plagiator</i> (Nees)	<i>Dendrocerus carpenteri</i> Curtis
<i>Eruca sativa</i> Mill.	<i>Myzus persicae</i> (Sulzer)	<i>Aphidius matricariae</i> Haliday	
		<i>Diaeretiella rapae</i> M'Intosh	
		<i>Ephedrus plagiator</i> (Nees)	
<i>Lactuca sativa</i> L.	<i>Uroleucon cichorii</i> (Koch)	<i>Ephedrus plagiator</i> (Nees)	
		<i>Aphidius matricariae</i> Haliday	
		<i>Monoctonus crepidis</i> (Haliday)	
<i>Petroselinum crispum</i> (P. Mill.) Nyman ex A. W. Hill	<i>Dysaphis</i> sp.	<i>Aphidius salicis</i> Haliday	
		<i>Myzus persicae</i> (Sulzer)	<i>Aphidius matricariae</i> Haliday
			<i>Ephedrus plagiator</i> (Nees)
			<i>Praon abjectum</i> (Haliday)
<i>Phaseolus vulgaris</i> L.	<i>Aphis fabae</i> Scopoli	<i>Aphidius ervi</i> Haliday	<i>Alloxysta brevis</i> Thomson
			<i>Lysiphlebus fabarum</i> (Marshall)
			<i>Pachyneuron aphidis</i> Bouché
			<i>Syrphophagus aphidivorus</i> Mayr
<i>Pisum sativum</i> L.	<i>Acyrtosiphon pisum</i> (Harris)	<i>Aphidius eadyi</i> Stary, Gonzalez & Hall	<i>Asaphes suspensus</i> Nees
			<i>Asaphes vulgaris</i> Walker
			<i>Pachyneuron aphidis</i> Bouché
			<i>Pachyneuron formosum</i> Walker
			<i>Praon barbatum</i> Mackauer
			<i>Praon volucre</i> (Haliday)
<i>Solanum tuberosum</i> L.	<i>Aphis gossypii</i> Glover	<i>Aphidius ervi</i> Haliday	<i>Asaphes suspensus</i> Nees
			<i>Dendrocerus aphidium</i> Rondani
			<i>Dendrocerus carpenteri</i> Curtis
<i>Valerianella locusta</i> L.	<i>Myzus persicae</i> (Sulzer)	<i>Lipolexis gracilis</i> (Forster)	
		<i>Lysiphlebus fabarum</i> (Marshall)	
		<i>Aphidius matricariae</i> Haliday	

Review of Interactions Between Host Plants, Aphids, Primary Parasitoids

Table 2. Review of trophic associations (host plant - host aphid - primary parasitoid - secondary parasitoid/hyperparasitoid) on some cereal crop plants in Slovenia.

Host plant	Host aphid	Primary parasitoid	Secondary parasitoid
<i>Avena sativa</i> L.	<i>Sitobion avenae</i> (F.)	<i>Aphidius ervi</i> Haliday	<i>Alloxysta brachyptera</i> Hartig
		<i>Aphidius rhopalosiph</i> de Stefani-Perez	<i>Alloxysta brevis</i> Thomson
		<i>Aphidius uzbekistanicus</i> Luzhetski	<i>Alloxysta victrix</i> Westwood
		<i>Ephedrus plagiator</i> (Nees)	<i>Dendrocerus carpenteri</i> Curtis
		<i>Praon volucre</i> (Haliday)	<i>Pachyneuron aphidis</i> Bouché
			<i>Pachyneuron muscarum</i> L.
			<i>Phaenoglyphis villosa</i> Hartig
<i>Hordeum vulgare</i> L.	<i>Sitobion avenae</i> (F.)	<i>Aphidius avenae</i> Haliday	<i>Alloxysta brevis</i> Thomson
	<i>Rhopalosiphum</i> sp.	<i>Aphidius ervi</i> Haliday	<i>Alloxysta victrix</i> Westwood
		<i>Aphidius rhopalosiph</i> de Stefani-Perez	<i>Asaphes suspensus</i> Nees
		<i>Aphidius uzbekistanicus</i> Luzhetski	<i>Asaphes vulgaris</i> Walker
		<i>Ephedrus plagiator</i> (Nees)	<i>Dendrocerus carpenteri</i> Curtis
		<i>Trioxys auctus</i> (Haliday)	<i>Phaenoglyphis villosa</i> Hartig
			<i>Syrphophagus aphidivorus</i> Mayr
<i>Triticum aestivum</i> L. emend. Fiori et Paol.	<i>Sitobion avenae</i> (F.)	<i>Aphidius avenae</i> Haliday	<i>Alloxysta brevis</i> Thomson
		<i>Aphidius ervi</i> Haliday	<i>Alloxysta victrix</i> Westwood
		<i>Aphidius rhopalosiph</i> de Stefani-Perez	<i>Asaphes suspensus</i> Nees
		<i>Aphidius uzbekistanicus</i> Luzhetski	<i>Asaphes vulgaris</i> Walker
		<i>Ephedrus plagiator</i> (Nees)	<i>Dendrocerus carpenteri</i> Curtis
		<i>Praon</i> sp.	<i>Dendrocerus aphidum</i> Rondani
			<i>Pachyneuron aphidis</i> Bouché
			<i>Pachyneuron muscarum</i> L.
			<i>Phaenoglyphis villosa</i> Hartig
<i>Zea mays</i> L.	<i>Rhopalosiphum maidis</i> (Fitch)	<i>Aphidius ervi</i> Haliday	<i>Alloxysta brevis</i> Thomson
	<i>Rhopalosiphum padi</i> (L.)	<i>Aphidius rhopalosiph</i> de Stefani-Perez	<i>Alloxysta victrix</i> Westwood
	<i>Sitobion avenae</i> (F.)	<i>Aphidius uzbekistanicus</i> Luzhetski	<i>Asaphes suspensus</i> Nees
		<i>Aphidius</i> sp.	<i>Asaphes vulgaris</i> Walker
		<i>Ephedrus plagiator</i> (Nees)	<i>Dendrocerus carpenteri</i> Curtis
		<i>Praon abjectum</i> (Haliday)	<i>Pachyneuron aphidis</i> Bouché
		<i>Praon</i> sp.	<i>Pachyneuron muscarum</i> L.
			<i>Phaenoglyphis villosa</i> Hartig
			<i>Syrphophagus aphidivorus</i> Mayr




Fig. 1. Relations of four trophic levels (secondary parasitoid/hyperparasitoid - host plant - host aphid - primary parasitoid) on four cereal crop plants in Slovenia collected from 2006-2010.

Table 3. Hyperparasitoids in vegetable and cereal crop plants in Slovenia.

	Vegetable crops			Cereal crops			Total (%)
	Female	Male	SUM	Female	Male	SUM	
<i>Alloxysta brachyptera</i>	/	/	0	/	1	1	1 (0,41%)
<i>Alloxysta brevis</i>	13	1	14	9	4	13	27 (11,0%)
<i>Alloxysta fulviceps</i>	1	/	1	/	/	0	1 (0,41%)
<i>Alloxysta victrix</i>	11	5	16	23	6	29	45 (18,3%)
<i>Asaphes suspensus</i>	3	3	6	19	10	29	35 (14,2%)
<i>Asaphes vulgaris</i>	3	1	4	4	9	13	17 (6,9%)
<i>Coruna clavata</i>	/	1	1	/	/	0	1 (0,41%)
<i>Dendrocerus aphidum</i>	1	1	2	1	/	1	3 (1,2%)
<i>Dendrocerus carpenteri</i>	3	2	5	20	11	31	36 (14,6%)
<i>Dendrocerus laticeps</i>	1	/	1	/	/	0	1 (0,41%)
<i>Pachyneuron aphidis</i>	5	4	9	14	10	24	33 (13,4%)
<i>Pachyneuron formosum</i>	1	/	1	/	/	0	1 (0,41%)
<i>Pachyneuron muscarum</i>	1	/	1	2	1	3	4 (1,63%)
<i>Phaenoglyphis villosa</i>	/	/	0	7	6	13	13 (5,3%)
<i>Syrphophagus aphidivorus</i>	18	8	26	/	2	2	28 (11,4%)
Total	61	26	87	99	60	159	246

DISCUSSION

Our survey on primary parasitoids in cereal crops reveals that *Aphidius ervi*, *A. rhopalosiphii*, *A. uzbekistanicus* and *E. plagiator* are the key species in the cereal aphid populations control in Slovenia. Other species of cereal parasitoids, among which also *A. avenae* Haliday, *P. volucre* Haliday, *Trioxys auctus* Haliday were found only occasionally in our study. These results are in agreement with findings in Germany (Höller *et al.*, 1993), Czech Republic (Stary, 1972, 1981), Poland (Pankanin-Franczyk and Sobota, 1998), Serbia (Tomanović *et al.*, 2008) and Denmark (Sigsgaard, 2002).

An analysis of trophic associations in vegetable crops reveals several key-stone species which should be considered in ecologically friendly management. *L. fabarum* can be used in sugar beet and pepper, *D. rapae* in cabbage and other vegetables from the family Brassicaceae, also *A. matricariae* can be introduced in pepper crop, and *A. salicis* Haliday on Apiaceae plants. Similarly specific assemblages of aphidiine parasitoids in vegetable crops were determined in other European countries (Kavallieratos *et al.*, 2004; Stary and Havelka, 2008; Tomanović and Brajković, 2001). *A. matricariae* and *L. fabarum* are rather eurytopic species which occur in steppe habitats including various vegetable crops, with exceptionally broad host range pattern in Southeast Europe (Kavallieratos *et al.*, 2004).

Primary aphid parasitoids are attacked by species-rich community of hyperparasitoids (Sullivan, 1988; Sullivan and Völkl, 1999) with different life-history strategies. *Alloxysta* spp. and *Phaenoglyphis* spp. develop as koinobiont endohyperparasitoids and parasitize the primary parasitoid larva within the living aphid. *Syrphophagus aphidivorus* develops also as koinobiont endohyperparasitoid, but parasitoid larvae can be attacked within living aphid and also inside aphid mummies. *Dendrocerus* spp. and all pteromalid species develop as idiobiont ectohyperparasitoids, where parasitoid prepupae and pupae inside aphid mummies are attacked (Sullivan, 1988; Sullivan and Völkl, 1999). Generally, externally feeding idiobiont ectohyperparasitoid species need less physiological adaptations for survival on a living host than koinobiont endohyperparasitoid species do (cit. by Sullivan and Völkl, 1999). All idiobiont hyperparasitoid species found in Slovenia belonging to genera *Dendrocerus*, *Asaphes*, *Pachyneuron* and *Coruna*, have a very broad range of hosts and attack various Aphidiinae genera and species, independent of the aphid host. Generally aphid endohyperparasitoid species are more numerous, but there are only a few species, namely *Alloxysta victrix* and *Phaenoglyphis villosa*, attacking a broad range of primary parasitoid hosts (Sullivan, 1988; Sullivan and Völkl, 1999).

In total, 22 primary parasitoid species and 15 species of hyperparasitoids in cereal and vegetable crops reveal a relatively diverse aphid parasitoid fauna on such small area, but that corresponds to diverse habitat, landscape and vegetation complexity in Slovenia. Increasing interest on biological control in Slovenia has led to improve the knowledge about four-trophic interactions among plants, aphids, aphidiine primary parasitoids and hyperparasitoids in crop plants.

ACKNOWLEDGEMENTS

We would like to thank to Aleksandar Stojanović (Belgrade Natural History Museum) for help in aphids and aphelinids identification. This research was partially supported within Professional Tasks from the Field of Plant Protection, a program funded by the Ministry of Agriculture, Forestry, and Food of Phytosanitary Administration of the Republic of Slovenia and within program III43001 (The Ministry of Education, Science and Technological development of the Republic of Serbia).

REFERENCES

- Agarwala, B. K., Das, J., 2012, Weed host specificity of the aphid, *Aphis spiraeicola*: developmental and reproductive performance of aphids in relation to plant growth and leaf chemicals of the Siam weed, *Chromolaena odorata*. *Journal of Insect Science*, 12(24): 1-13.
- Bianchi, F. J., Wäckers, F. L., 2008, Effects of flower attractiveness and nectar availability in field margins on biological control by parasitoids. *Biological control*, 55(6): 719-727.
- Blande, J. D., Pickett, J. A., Poppy, G. M., 2007, A Comparison of Semiochemically Mediated Interactions Involving Specialist and Generalist *Brassica*-feeding Aphids and the Braconid Parasitoid *Diaeretiella rapae*. *Journal of Chemical Ecology*, 33: 767-779.
- Boller, E. F., van Lenteren, J. C., Delucchi, V., 2006, *International Organization for Biological Control of Noxious Animals and Plants - History of the first 50 years (1956-2006)*. IOBC, OILB, Ponsen & Looijen, Wageningen, The Netherlands, 275.
- Boivin, G., Brodeur, J., 2006, *Intra- and Interspecific Interactions among Parasitoids: Mechanisms, Outcomes and Biological Control*. In: Brodeur, J., Boivin, G. (Eds.) *Trophic and Guild Interactions in Biological Control*. Springer, 123-144.
- Bosch, R., van den Telford, A. D., 1964, *Environmental modification and biological control*. In: DeBac, P. (Ed.) *Biological control of Insect Pests and Weeds*. New York, Reinhold, 459-488.
- Dicke, M., Sabelis, M. W., 1988, How plants obtain predatory mites as bodyguards. *Netherlands Journal of Zoology*, 38: 148-165.
- Godfray, H. C. J., 1994, *Parasitoids. Behavior and Evolutionary Ecology*. Princeton, New Jersey: Princeton University Press, 488.
- Henry, L. M., May, N., Acheampong, S., Gillespie, D. R., Roitberg, B. D., 2010, Host-adapted parasitoids in biological control: Does source matter? *Ecological Applications*, 20: 242-250.
- Höller, C., Borgemeister, C., Haardt, H., Powell, W., 1993, The Relationship between Primary Parasitoids and Hyperparasitoids of Cereal Aphids: An Analysis of Field Data. *Journal of Animal Ecology*, 62(1): 12-21.
- Kavallieratos, N. G., Lykouressis, P., Sarlis, G. P., Stathas, G. J., Sanchís-Segovia, A., Athanassiou, C. G., 2001, The Aphidiinae (Hymenoptera: Ichneumonoidea: Braconidae) of Greece. *Phytoparasitica*, 29: 306-340.
- Kavallieratos, N. G., Tomanović, Ž., Starý, P., Athanassiou, C. G., Sarlis, G. P., Petrović, O., Niketić, M., Veroniki, M. A., 2004, A survey of aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) of southeastern Europe and their aphid-plant associations. *Applied Entomology and Zoology*, 39: 527-563.
- Kavallieratos, N. G., Tomanović, Ž., Starý, P., Athanassiou, C. G., Fasseas, C., Petrović, O., Stanisavljević, L., Anagnouveroniki, M., 2005, *Praon* Haliday (Hymenoptera: Braconidae: Aphidiinae) of Southeastern Europe: key, host range and phylogenetic relationship. *Zoologische Anzeiger*, 243: 181-209.
- Kos, K., Tomanović, Ž., Petrović-Obradović, O., Laznik, Ž., Vidrih, M., Trdan, S., 2008, Aphids (Aphididae) and their parasitoids in selected vegetable ecosystems in Slovenia. *Acta agriculturae Slovenica*, 91 (1): 15-22.
- Kos, K., Tomanović, Ž., Rojht, H., Trdan, S., 2009, Aphid parasitoids in Slovenia and their significance in biological control. *Acta entomologica Slovenica*, 17(1): 5-12 [in Slovene].

Review of Interactions Between Host Plants, Aphids, Primary Parasitoids

- Kos, K., Trdan, S., Petrović, A., Starý, P., Kavallieratos, N., Petrović-Obradović, O., Tomanović, Ž., 2012, Aphidiinae (Hymenoptera, Braconidae, Aphidiinae) from Slovenia, with description of a new Aphidius species. *Zootaxa*, 3456: 36-50.
- Landis, D. A., Wratten, S. D., Gurr, G. M., 2000, Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology*, 45: 175-201.
- Micha, S. G., Kistenmacher, S., Mölock, G., Wyss, U., 2000, Tritrophic interactions between cereals, aphids and parasitoids: Discrimination of different plant-host complexes by *Aphidius rhopalosiphii* (Hymenoptera: Aphidiidae). *European Journal of Entomology*, 97: 539-543.
- Milevoj, L., 2001, The role of native beneficial organisms in biological control. *Lectures and papers of 5th Slovenian conference on plant protection, 6.-8. of March 2001, Čatež ob Savi*. Ljubljana, Plant Protection Society of Slovenia, Slovenia: 59-63 [in Slovene].
- Minks, A. K., Harrewijn P., 1988, *Aphids, their biology, natural enemies and control*. World Crop Pests 2B. Amsterdam, Elsevier, 364.
- Modic, Š., Urek, G., 2008, Contribution to the knowledge of the aphid fauna (Sternorrhyncha: Aphidoidea) of Slovenia. *Acta Entomologica Slovenica*, 16(1): 87-97 [in Slovene].
- Pankanin Franczyk, M., Sobota, G., 1998, Relationships between primary and secondary parasitoids of cereal aphids. *Journal of Applied Entomology*, 122(7): 389-395.
- Parker, W. E., Howard, J. J., Foster, S. P., Denholm, I., 2006, The effect of insecticide application sequences on the control and insecticide resistance status of the peach-potato aphid, *Myzus persicae* (Hemiptera : Aphididae), on field crops of potato. *Pest Management Science*, 62(4): 307-315.
- Price, P. W., 1984, *Insect Ecology*. 2nd ed., Wiley, New York, 607.
- Quacchia, A., Moriya, S., Bosio, G., Scapin, I., Alma, A., 2008, Rearing, release and settlement prospect in Italy of *Torymus sinensis*, the biological control agent of the chestnut gall wasp *Dryocosmus kuriphilus*. *BioControl*, 53: 829-839.
- Sigsgaard, L., 2002, A survey of aphids and aphid parasitoids in cereal fields in Denmark, and the parasitoids' role in biological control. *Journal of Applied Entomology*, 126: 101-107.
- Starý, P., 1970, *Biology of Aphid Parasites, with Respect to Integrated Control*. Series Entomologica 6. Junk Publishers, Hague, The Netherlands, 651.
- Starý, P., 1972, *Aphidius uzbekistanicus* Luzhetskii (Hym., Aphidiidae) a parasite of graminicolous pest aphids. *Annotationes Zoologicae et Botanicae. Bratislava*, 85: 1-7.
- Starý, P., 1981, Biosystematic synopsis of parasitoids on cereal aphids in the western Palearctic (Hymenoptera, Aphidiidae; Homoptera, Aphidoidea). *Acta Entomologica Bohemoslovaca*, 78: 382-396.
- Starý, P., 1988, *Parasites: Aphidiidae*. In: Minks, A. K., Harrewijn, P. (Eds.) *Aphids, their biology, natural enemies and control*. World Crop Pests 2B. Amsterdam, Elsevier, 171-184.
- Starý, P., Havelka, J., 2008, Fauna and associations of aphid parasitoids in an up-dated farmland area (Czech Republic). *Bulletin of Insectology*, 61: 251-276.
- Stoerck, A., Poppy, G. M., van Emden, H. F., Powell, W., 2000, The role of plant chemical cues in determining host preference in the generalist aphid parasitoid *Aphidius colemani*. *Entomologia Experimentalis et Applicata*, 97: 41-46.
- Strand, M. R., 2000, *Developmental Traits and Life-History Evolution in Parasitoids*. In: Hochberg, M. E., Ives, A. R. (Eds.) *Parasitoid population Biology*. Princeton, New Jersey, Princeton University Press, 139-162.
- Sullivan, D. J., 1988, *Aphid hyperparasites*. In: Minks, A. K., Harrewijn, P. (Eds.) *Aphids, Their Biology, Natural Enemies and Control*, World Crop Pests 2B Elsevier, Amsterdam, 189-203.
- Sullivan, D. J., Völkl, W., 1999, Hyperparasitism: Multitrophic Ecology and Behavior. *Annual Review of Entomology*, 44: 291-315.
- Tomanović, Ž., Brajković, M., 2001, Aphid parasitoids (Hymenoptera, Aphidiidae) of agroecosystems of the south part of the Pannonian area. *Archives of Biological Science Belgrade*, 53, 57-64.

- Tomanović, Ž., Kavallieratos, N. G., Starý, P., Petrović-Obradović, O., Athanassiou, C., Stanislavljević, L. Ž., 2008, Cereal aphids (Hemiptera: Aphidoidea) in Serbia: Seasonal dynamics and natural enemies. *European Journal of Entomology*, 105: 195-501.
- Tscharntke, T., Hawkins, B. A., 2002, *Plant genetic variation in tritrophic interactions*. In: Tscharntke, T., Hawkins, B. A. (Eds.) *Multitrophic Level Interactions*. Cambridge, Cambridge University Press, 8-43.
- Turlings, T. C. J., Tumlinson, J. H., Lewis, W. J., 1990, Exploitation of Herbivore-Induced Plant Odors by Host-Seeking Parasitic Wasps. *Science*, 250: 1251-1253.
- Zuñiga, E., 1986, Control biológico de los áfidos (Hom.; Aphididae) de los cereales en Chile. II Obtención, introducción y cuarentena de depredadores y parasitoides. *Agricultura Técnica (Santiago)*, 46(4): 479-487.
- Wajnberg, E., 2010, *Genetics of the behavioral ecology of egg parasitoids*. In: Consóli F. L., Parra, J. R. P., Zucchi, R.A. (Eds.) *Egg Parasitoids in Agroecosystems with Emphasis on Trichogramma*. Springer, 149-165.

Received: February 24, 2011

Accepted: November 08, 2012