

## New Records on the Ichneumonidae Fauna (Hymenoptera) of the Black Sea Coast of Turkey

Janko KOLAROV<sup>1</sup>      Saliha ÇORUH<sup>2\*</sup>

<sup>1</sup>Faculty of Pedagogy, University of Plovdiv, Plovdiv, BULGARIA

<sup>2</sup>Atatürk University, Faculty of Agriculture, Department of Plant Protection, 25240 Erzurum, TURKEY

e-mails: <sup>1</sup>j.kolarov@abv.bg, <sup>2</sup>\*spekel@atauni.edu.tr

ORCID IDs: <sup>1</sup>0000-0002-0027-4162, <sup>2</sup>0000-0002-6822-6677

### ABSTRACT

In this study, Ichneumonidae (Hymenoptera) species collected from Black Sea Coast (Rize) of Turkey are presented. In total 15 species belonging to the subfamilies Metopiinae, Ophioninae, Orthocentrinae, Oxytorinae, Pimplinae and Tryphoninae were determined. Of these, *Exochus alpinus* (Zetterstedt, 1838), *Cylloceria borealis* (Roman, 1925), *Orthocentrus patulus* Holmgren, 1858 and *Zatypota albicoxa* (Walker, 1874) are new records for the Turkish fauna. In addition, new localities for the other species are recorded. We discuss the geographical and zoogeographical records of all the species.

**Key words:** Darwin wasps, Hymenoptera, Ichneumonidae, new records, Rize, Turkey.

## INTRODUCTION

Hymenoptera is one of the four large insect orders exceeding 100,000 species in the world, the other major orders being Coleoptera, Lepidoptera and Diptera (Gauld & Bolton, 1988; Wahl & Micheal, 1993).

Ecologically and economically few groups of insects are so important to mankind as the Hymenoptera. Bees provide the vital ecosystem service of pollination in both natural and managed systems (Gallai, Salles, Settele, & Vaissiere, 2009) while parasitic Hymenoptera control populations of phytophagous insects (Tscharntke, Bommarco, Clough, Crist, & Kleijn, 2007) and can be effective agents for control of pest insects (Bale, van Lenteren, & Bigler, 2008; Brodeur & Boivin 2004; Jonsson, Wratten, Landis, & Gurr, 2008).

Ichneumonidae is the first megadiverse Apocrita family in the world with 25,285 species (Yu, Achterberg, & Horstmann, 2016). The family is divided into more than 40 subfamilies (Broud, Shaw, & Fitton, 2018). Consequently, the biology of ichneumonids is extremely diverse. Ichneumonids mostly parasitize the immature stages of the Holometabola, and are frequently associated with Lepidoptera and sawflies (Hymenoptera). Ectoparasitism is considered the primitive condition and endoparasitism has evolved several times independently within the family (Rasplus, Villemant, Paiva, Delvare, & Roques, 2010).

Studies on Ichneumonidae of Turkey have gained acceleration in recent years. Çoruh, Gürbüz, Kolarov, Yurtcan, & Özdan (2013), reported 975 species in 282 genera for Turkey Ichneumonidae fauna. Now, the number of Ichneumonidae species has reached 1316 with valuable studies: Yaman, 2014; Özdan, 2014; Çoruh, Kolarov, & Çoruh, 2014a; Çoruh, Kolarov, & Özbek, 2014b; Kolarov, Çoruh, & Çoruh, 2014a; Kolarov, Yıldırım, Çoruh & Yüksel, 2014b; Riedel, Kolarov, Çoruh & Çoruh, 2014; Kolarov, Çoruh, & Çoruh, 2015; Kolarov, Çoruh, & Çoruh, 2016; Çoruh, 2017; Kolarov, Çoruh, & Çoruh, 2017; Narmanlioğlu & Çoruh, 2017; Çoruh, Kolarov, & Çoruh, 2018; Riedel, Diller, & Çoruh, 2018; Sarı & Çoruh, 2018; Çoruh, 2019a, b; Vas, 2019a, b; Özdan & Gürbüz, 2019; Çoruh, Kolarov & Ercelep, 2019; Çaylak & Çoruh 2020a, b; Kolarov, Çoruh, & Çoruh, 2020; Kıracı & Gürbüz, 2020; Schwarz, 2020, Teymuroğlu, & Çoruh 2021, Kolarov, Çoruh & Ercelep, 2021, Yurtcan, Çoruh, Kolarov, Özdan, Gürbüz, & Erkaya, 2021; Çoruh, Kolarov, & Ercelep, 2022).

The purpose of this study is to identify the Ichneumonidae species collected from Black sea region, to make this data available to researchers and relevant people and to contribute the biodiversity.

## MATERIAL AND METHODS

### Data sampling

Adult ichneumonid samples were collected by sweeping net on flowering plants in six different localities and altitudes at above level (10-200 m) (Table 1) of Rize (Fig. 1, 2) in Turkey (Anatolia), during summer of 2016-2017. Collected specimens were preserved in 70 % ethanol. All examined material was collected by the second

### New Records on the Ichneumonidae Fauna

author and Halil Çoruh and determined by authors and deposited in the Entomology Museum Erzurum, Turkey (EMET). New record of species is marked by an asterisk (\*). General distributions (Table. 2), hosts and associated plants of the species were taken from Yu et al, (2016).

Table 1. Data of collected species.

	Locality	Year	Altitude (M.)
Rize	Ardeşen	2016-2017	10
	Çayeli		10
	Derepazari		50
	Güneysu		150
	İyidere		210
	Pazar		37

Table 2. Zoogeographic distribution of collected species.

Zoogeographical Regions	Afrotropical	Eastern Palearctic	European Region	Nearctic	Neotropical	Oceanic	Oriental	Western Palearctic
<i>Colpotrochia cincta</i>	•	•						•
<i>Exochus alpinus</i>				•				
<i>Exochus prosopius</i>	•	•						•
<i>Enicospilus ramidulus</i>	•	•	•				•	•
<i>Cyllooceria borealis</i>		•	•				•	•
<i>Orthocentrus patulus</i>			•					•
<i>Oxytorus luridator</i>			•					•
<i>Clistopyga rufator</i>		•	•					•
<i>Pimpla aquilonia</i>		•	•	•				•
<i>Pimpla rufipes</i>			•		•			•
<i>Pimpla spuria</i>		•	•				•	•
<i>Scambus nigricans</i>		•	•					•
<i>Zatypota albicoxa</i>		•	•				•	•
<i>Ctenochira angulata</i>		•	•					•
<i>Oedemopsis scabricula</i>		•	•	•			•	•



Fig. 1. Map of study area.



Fig 2. Photographs of study area.

### Study area

Rize is a province located in the northeastern Turkey and has a coast to the Black Sea. Located in the Black Sea Region, Rize is surrounded by Trabzon in the west, Artvin in the east, Bayburt in the southwest, and Erzurum in the south (Anonymous, 2021).

In general, the land of Rize province are covered with a very diverse and very lush vegetation. Nearly 48% of Rize provincial lands are covered with forest and heaths. Forest are found up to an altitude of 2800 m. 24% is covered with cultivated and plated areas.

## RESULTS

During the investigation, we found 15 species belonging to Metopiinae, Ophioninae Ophioninae, Orthocentrinae, Oxytorinae, Pimplinae and Tryphoninae. The species list is given below.

### Subfamily Metopiinae Förster, 1869

#### *Colpotrochia cincta* (Scopoli, 1763)

Material examined: Rize: İyidere, 210 m, 23.07.2016, 2 ♀♀; 02.08.2017, 1 ♂.

Distribution: Palaearctic, known from Turkey (Kolarov, Çoruh, Yurtcan, & Gürbüz, 2009; Kolarov & Çalışmaşır, 2011; Çoruh & Özbeğ, 2011; Çoruh, Kolarov, & Çoruh, 2014a; Çoruh, Kolarov, & Özbeğ, 2014b; Kolarov, Yıldırım, Çoruh, & Yüksel, 2014b).

Hosts: *Acronicta aceris*, *Acronicta rumicis*, *Aletia conigera*, *Axylia*, *Mythimna turca*, *Papestra biren*.

Associate plant: *Peucedanum oreoselinum*.

*New Records on the Ichneumonidae Fauna*

**\**Exochus alpinus* (Zetterstedt, 1838)**

Material examined: Rize: Ardeşen, 02.08.2017, 2 ♀♀; Güneysu, 150 m, 16.06.2016, 1 ♀.

Distribution: Palaearctic, new for Turkey.

Hosts: *Acleris rhombana*, *Apotomis sauciana*, *Archips xylosteana*, *Epinotia solandriana*, *Epinotia tetraquetra*, *Hedya dimidiana*, *Phauda flammans*, *Triaxomera parasitella*.

***Exochus prosopius* Gravenhorst, 1829**

Material examined: Rize: İyidere, 210 m, 29.06.2016, 1 ♀; 23.07.2017, 2 ♂♂; 02.08.2017, 1 ♂.

Distribution: Palaearctic, known from Turkey (Kolarov, Çoruh, Yurtcan & Gürbüz, 2009; Çoruh et al, 2014b; Kolarov, Çoruh, & Çoruh 2017).

**Subfamily Ophioninae Shuckard, 1840**

***Enicospilus ramidulus* (Linnaeus, 1758)**

Material examined: Rize: Güneysu, 150 m, 05.07.2017, 1 ♀; İyidere, 210 m, 09.06.2017, 2 ♂♂ 5 ♀♀.

Distribution: Afrotropical, Palaearctic, Oriental, known from Turkey (Kolarov, 1995; Kolarov, Pekel, Özbek, Yıldırım, & Çalmaşur, 2000; Gürbüz, 2004, 2005; Beyarslan, Yurtcan, Erdoğan, & Aydoğdu, 2006; Okyar & Yurtcan, 2007; Çoruh & Çoruh, 2008, Gürbüz, Kirtay & Birol, 2009; Kıracı & Gürbüz, 2020; Çoruh et al, 2014b; Kolarov, Çoruh & Çoruh, 2014a; Çoruh & Çalmaşur, 2016; Kolarov et al, 2017; Kolarov et al, 2020).

Hosts: *Agrotis segetum*, *Anarta myrtillii*, *Axylia putris* [*Nicotiana tabacum*], *Catocala electa*, *Cucullia absinthii*, *Dendrolimus pini*, *Dypterygia scabriuscula*, *Eriogaster lanestris*, *Hadena bicruris*, *Hadena rivularis*, *Hyles euphorbiae*, *Hyphantria cunea* [*Corylus avellana*], *Lacanobia oleracea*, *L. pisi*, *Melanchra persicariae*, *Nematus salicis*, *Ochropleura praecox*, *Panolis flammea*, *Phalera bucephala*, *Phlogophora meticulosa*, *Poecilocampa populi*.

Associate plants: *Alnus glutinosa*, *Carum carvi*, *Oryza sativa*, *Salvia glutinosa*, *Seseli libanotis*.

**Subfamily Orthocentrinae Förster, 1869**

**\**Cylloceria borealis* (Roman, 1925)**

Material examined: Rize: İyidere, 210 m, 09.06.2017, 1 ♂.

Distribution: Holarctic, new for Turkey.

**\**Orthocentrus patulus* Holmgren, 1858**

Material examined: Rize: Pazar, 37 m, 31.08.2017, 1 ♀.

Distribution: Europe, new for Turkey.

### **Subfamily Oxytorinae Thomson, 1883**

#### ***Oxytorus luridator* (Gravenhorst, 1820)**

Material examined: Rize: Pazar, 37 m, 31.08.2017, 1 ♀.

Distribution: Europe, known for Turkey (Kolarov, Çoruh & Çoruh, 2015; Kolarov, Çoruh & Çoruh, 2016).

Associate plants: *Sparganium* sp.

### **Subfamily Pimplinae**

#### ***Clistopyga rufator* Holmgren, 1856**

Material examined: Rize: Çayeli, 10 m, 16.06.2017, 1 ♀.

Distribution: Palaearctic, known from Turkey (Yurtcan, 2004, 2007; Çoruh, 2005; Çoruh, Özbek, Kolarov, 2007; Buncukçu, 2008; Çoruh & Özbek, 2008; Gürbüz, Ljubomirov, Kolarov, Yurtcan, Tabur, Çoruh & Buncukçu, 2008; Gürbüz et al, 2009; Çoruh, 2009, 2010; Birol, 2010; Çoruh & Kolarov, 2010; Çoruh, Çalmaşur & Tozlu, 2010; Özdan, 2014; Çoruh et al, 2014b; Çoruh, 2016; et al, 2021).

Host: *Clubiona juvenis*.

#### ***Pimpla aquilonia* Cresson, 1870**

Material examined: Rize: İyidere, 210 m, 09.06.2017, 1 ♂; Pazar, 37 m, 11.07.2017, 1 ♂.

Distribution: Holarctic, known from Turkey (Yurtcan & Beyarslan, 2005, Çoruh & Kolarov, 2010; Çoruh et al, 2014a; Çoruh, 2016)

Hosts: *Archips fuscocupreanus*, *Argiope argentata*, *Choristoneura diversana*, *Coleophora serratella*, *Croesia semipurpurana*, *Cymolomia hartigiana*, *Dendrolimus alboleatus*, *Epinotia nigricana*, *Grapholita molesta*, *Lambdina fiscellaria* [*Abies balsamea*], *Lambdina fiscellaria lugubrosa*, *Magdalisa frontalis*, *Orgyia leucostigma*, *Orthosia hibisci*, *Parasteatoda tepidariorum*, *Psilocorsis cryptolechiella*, *Rheumaptera hastata*, *Spilonota ocellana*, *Teleiodes saltuum*, *Tortrix viridana*, *Zeiraphera griseana*,

Associate plants: *Listera ovata*.

#### ***Pimpla rufipes* Brullé, 1846**

Material examined: Rize: İyidere, 210 m, 23.06.2017, 1 ♀.

Distribution: Palaearctic, Neotropical, known from Turkey (Çoruh, 2005, 2009; Okyar & Yurtcan, 2007; Çoruh & Özbek, 2008; Kolarov & Çalmaşur, 2011; Özbek & Çoruh, 2012, Çoruh, 2016).

Hosts: *Hyphantria cunea*, *Malacosoma neustria*, *Plusia* sp.

Remarks: This species obtained from *Malacosoma neustria* and *M. franconica*, *Yponomeuta malinella* in our previous study (Çoruh & Özbek, 2008).

New Records on the Ichneumonidae Fauna

**Pimpla spuria Gravenhorst, 1829**

Material examined: Rize: Derepazarı, 50 m, 23.06.2016, 1 ♂; Pazar, 37 m, 04.07.2017, 1 ♂.

Distribution: Palaearctic, Oriental, known for Turkey (Fahringer, 1922; Özdemir & Kılınçer, 1990; Öncüler, 1991; Kolarov & Beyarslan, 1994; Kolarov, Yurtcan, & Beyarslan, 1997a; Kolarov, Beyarslan & Yurtcan, 1997 b; Kolarov, Özbek, & Yıldırım, 1999; Kolarov, Yurtcan & Beyarslan, 2002; Gürbüz, 2005; Kolarov & Gürbüz, 2004; Çoruh, 2005; Gürbüz, 2005; Yurtcan & Beyarslan, 2005; Kirtay, 2008; Çoruh, Çalışmaz, & Tozlu, 2010; Çoruh & Kolarov, 2010; Eroğlu, Kıraç & Birol, 2011; Kolarov et al, 2014b; Çoruh et al, 2014a; Çoruh, 2016; Kolarov et al, 2016; Sarı & Çoruh, 2018; Kıraç & Gürbüz, 2020).

Hosts: *Agonopterix heracliana*, *Agonopterix propinquella*, *Archips rosana*, *Calliteara pudibunda*, *Canephora hirsuta*, *Cydia funebrana*, *Cydia pomonella*, *Cynaeda dentalis*, *Depressaria pastinacella*, *Enarmonia formosana*, *Etiella zinckenella*, *Eupithecia vulgata*, *Eupoecilia ambiguella*, *Gypsonoma aceriana*, *Hyphantria cunea*, *Larentia clavaria*, *Lobesia botrana*, *Loxostege sticticalis*, *Lymantria dispar*, *Naranga aenescens*, *Ostrinia nubilalis*, *Pempelia genistella*, *Psoricoptera gibbosella*, *Sesamia nonagrioides*, *Tebenna bjerkandrella*, *Yponomeuta malinella*, *Yponomeuta padella*.

Associated plants: *Acer campestre*, *Anethum graveolens*, *Chaerophyllum bulbosum*, *Daucus carota*, *Daucus carota sativus*, *Euphorbia nicaeensis*, *Heracleum sphondylium*, *Tamarix* sp.

Remarks: This species obtained from *Malacosoma neustria* and *Yponomeuta malinella* in our previous study (Çoruh & Özbek, 2008).

**Scambus nigricans Thomson, 1877**

Material examined: İyidere, 210 m, 05.08.2017, 1 ♀; Pazar, 37 m, 23.07.2017, 1 ♂, 01.08.2017, 1 ♂.

Distribution: Palaearctic, known from Turkey (Kolarov & Beyarslan, 1994; Kolarov et al, 1997a; Kolarov et al, 1999; Kolarov et al, 2002; Kolarov & Gürbüz, 2004; Çoruh, 2005; Yurtcan, 2007; Çoruh et al, 2007, Çoruh & Kolarov, 2010; Çoruh et al, 2010; Kolarov & Çalışmaz, 2011; Özdek & Çoruh, 2012; Çoruh et al, 2014b; Çoruh, 2016; Kolarov et al, 2017; Kolarov et al, 2020; Yurtcan et al, 2021).

Hosts: *Adaina microdactyla*, *Aethes francillana*, *Aethes williana*, *Agonopterix heracliana*, *Alsophila aescularia*, *Apocheima hispidaria*, *Cnephasia stephensi*, *Cochylis atricapitana*, *Cydia delineana*, *Cydia discretana*, *Cydia pactolana*, *Depressaria daucella*, *Depressaria pastinacella*, *Dioryctria simplicella*, *Epiblema cirsiana*, *Epiblema scutulana*, *Gortyna flavago*, *Hartigia linearis*, *Heptamelus ochroleucus*, *Lampronia morosa*, *Lipara lucens*, *Lycaena dispar batava*, *Lycia pomonaria*, *Malacosoma neustria*, *Mompha conturbatella*, *Myelois circumvoluta*, *Noeeta pupillata*, *Oxyna parietina*, *Phigalia pilosaria*, *Platyptilia nemoralis*, *Pseudaletia unipuncta*, *Rhyacionia buoliana*, *Scrobipalpa clintoni*, *Triedris paleana*.

Associated plants: *Anethum graveolens*, *Chaerophyllum bulbosum*, *Daucus carota*, *Euphorbia nicaeensis*, *Heracleum sphondylium*, *Peucedanum oreoselinum*.

Remark: This species was collected on *Heracleum pastinacifolium* in our previous study (Kolarov et al, 2020).

### **\**Zatypota albicoxa* (Walker, 1874)**

Material examined: Rize: Pazar, 37 m, 19.07.2017, 2 ♀♀.

Distribution: Palaearctic, Oriental, new for Turkey.

Hosts: *Parasteatoda lunata*, *Parasteatoda merapiensis*, *Parasteatoda oculiprominens*, *Parasteatoda simulans*, *Parasteatoda tabulata*, *Parasteatoda tepidariorum*, *Theridion limatum*.

Associate plants: *Oryza sativa*.

### **Subfamily Tryphoninae Shuckard, 1840**

#### ***Ctenochira angulata* (Thomson, 1883)**

Material examined: Rize: Pazar, 37 m, 27.08.2017, 1 ♂.

Distribution: Palaearctic, known from Turkey (Yurtcan & Beyarslan, 2002; Yaman, 2014; Kolarov et al, 2016; Çoruh, 2019).

#### ***Oedemopsis scabricula* (Gravenhorst, 1829)**

Material examined: Rize: Ardeşen, 10 m, 05.08.2017, 3 ♀♀.

Distribution: Holarctic, Oriental, known from Turkey (Çoruh, Özbek, Kolarov, 2005; Beyarslan, Erdoğan, Çetin, & Aydoğdu, 2006; Çoruh et al, 2014a; 2014b; Yaman, 2014; Çoruh, 2019, Kolarov et al, 2017).

Hosts: *Acleris variegana*, *Aleimma loeflingiana*, *Ancylis apicella*, *Archips oporana*, *Archips rosana*, *Blastobasis lignea*, *Cacoecimorpha pronubana*, *Choristoneura rosaceana*, *Cladius difformis*, *Clepsis costana*, *Croesia holmiana*, *Ditula angustiorana*, *Epinotia albangulana*, *Epinotia solandriana*, *Epiphyas postvittana*, *Eudemis profundana*, *Euproctis similis*, *Pandemis cerasana*, *Pandemis heparana*, *Ptycholoma lecheana*, *Rhyacionia buoliana*, *Selania leplastriana*, *Spilonota ocellana*, *Tortrix viridana*, *Yponomeuta cagnagella*.

### **Zoogeographical notes**

*Enicospilus ramidulus* is the species with the multiregional ranges is the species *Enicospilus ramidulus*, distributed in Afrotropical, Palaearctic and Oriental zoogeographical regions.

Three species have distributional ranges in two zoogeographical regions of them: *Pimpla rufipes* distributed in Western Palaearctic and Neotropical regions; *Pimpla spuria* and *Zatypota albicoxa* distributed in Palaearctic and Oriental regions.

Two species, *Cylloceria borealis* and *Pimpla aquilonia* have Holarctic distribution. Most of the species have Palaearctic distribution. Of them, *Colpotrochia cincta*, *Exochus alpinus*, *E. prosopius*, *Clistopyga rufator*, *Scambus nigricans* and *Ctenochira angulata*.

European distribution ranges have 2 species - *Orthocentrus patulus* and *Oxytorus luridator*.

## REFERENCES

- Anonymous, 2021(September, 1), Rize. Retrieved from [https://tr.wikipedia.org/wiki/Rize\\_\(il\)](https://tr.wikipedia.org/wiki/Rize_(il))
- Bale, J. S., Van Lenteren, J. C. & Bigler, F. (2008). Biological control and sustainable food production. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 363, 761-776.
- Beyarslan, A., Yurtcan, M., Erdođan, Ö.Ç., & Aydođdu, M. (2006). A study on Braconidae and Ichneumonidae from Ganos Mountains (Thrace Region, Turkey) (Hymenoptera, Braconidae, Ichneumonidae). *Linzer Biologische Beiträge*, 38(1), 409-422.
- Birol, O. (2010). *Isparta İli Davraz Dağı Ichneumonidae (Hymenoptera) Faunası Üzerine Bir Araştırma*. Süleyman Demirel Üniversitesi, Fen Bilimleri Enstitüsü, Isparta, 71 s.
- Buncukçu, A., 2008, *Isparta İli Merkez ve Adana, Yumurtalık İlçesi-Halep Çamlığı Ichneumonidae Türlerinin Tespiti ve Kültüre Edilebilen Türlerin Biyolojilerinin Araştırılması*. Süleyman Demirel Üniversitesi, Fen Bilimleri Enstitüsü, Isparta, 74 s.
- Brodeur, J. & Boivin, G. (2004). Functional ecology of immature parasitoids. *Annual Review of Entomology*, 49, 27-49.
- Broud, G., Shaw, M., & Fitton, M. (2018). Ichneumonid Wasps (Hymenoptera: Ichneumonidae): their Classification and Biology. *Handbooks for the Identification of British Insects*, 7(12), 1-418.
- Çaylak, F.Z. & Çoruh, S. (2020a). First record of *Woldstedtius citropectoralis* Schmiedeknecht, 1926 (Hymenoptera: Ichneumonidae: Diplazoninae) from Turkey. *Munis Entomology & Zoology*, 15(2), 457-462.
- Çaylak, F.Z. & Çoruh, S. (2020b). Contribution to the knowledge of Ichneumonidae (Hymenoptera) of Bursa Uludağ National Park area including new records. *Turkish Journal of Entomology*, 43(4), 503-517.
- Çoruh, S. (2005). Erzurum ve Çevre İllerdeki Pimplinae (Hymenoptera, Ichneumonidae) Türleri Üzerinde Faunistik, Sistematiğ ve Ekolojik Çalışmalar. Atatürk Üniversitesi, Fen Bilimleri Enstitüsü, Erzurum, 212 s.
- Çoruh, S. (2009). Two little-known Ichneumonidae (Hymenoptera) from Turkey, including one new for the Turkish fauna. *Zoology in the Middle East*, 48: 106-107.
- Çoruh, S. (2010). Composition, habitat distribution and seasonal activity of Pimplinae (Hymenoptera: Ichneumonidae) in North-East Anatolia region of Turkey. *Anadolu Tarım Bilimleri Dergisi*, 25(1): 28-36.
- Çoruh, S. (2016). Biogeography and Host Evaluation of the Subfamily Pimplinae (Hymenoptera: Ichneumonidae) in Turkey. *Journal of the Entomological Research Society*, 18(2), 33-66.
- Çoruh, S. (2017). Taxonomical and biogeographical evaluation of the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae) in Turkey. *Entomofauna*, 38(21), 425-476.
- Çoruh, S. (2019a). Taxonomic and biogeographic evaluations of the subfamily Cryptinae (Hymenoptera: Ichneumonidae). *Turkish Journal of Entomology*, 43(3), 313-337.
- Çoruh, S. (2019b). Taxonomical and biogeographical evaluation of the subfamily Tryphoninae (Hymenoptera: Ichneumonidae) in Turkey. *Journal of the Entomological Research Society*, 21(3), 301-321.
- Çoruh, İ. & Çoruh, S. (2008). Ichneumonidae (Hymenoptera) species associated with some Umbelliferae plants occurring in Palandöken Mountains of Erzurum, Turkey. *Turkish Journal of Zoology*, 32(2), 121-124.
- Çoruh, S. & Özbek, H. (2008). New and rare Ichneumonidae (Hymenoptera) species from Turkey. *Zoology in the Middle East*, 43, 114-116.
- Çoruh, S. & Kolarov, J. (2010). A review of the Turkish Orthopelmatinae (Insecta: Hymenoptera: Ichneumonidae). *Scientific Research and Essays*, 5(22), 3518-3521.
- Çoruh, S. and Özbek, H. (2011). New and little known some Ichneumonidae (Hymenoptera) species from Turkey with some ecological notes. *Turkish Journal of Entomology*, 35(1), 119-131.
- Çoruh, S. & Çalmaşur, Ö. (2016). A new and additional records of the Ichneumonidae (Hymenoptera) from Turkey. *Turkish Journal of Zoology*, 40(4), 625-629.
- Çoruh, S., Özbek, H. & Kolarov, J. (2005). A contribution to the knowledge of Tryphoninae (Hymenoptera: Ichneumonidae) from Turkey. *Zoology in the Middle East*, 35: 93-98.

- Çoruh, S., Özbeğ, H., & Kolarov, J. (2007). Aras Vadisi (Kars)'ne yerleştirilen malezya tuzağından elde edilen Ichneumonidae (Hymenoptera) türleri. Türkiye II. Bitki Koruma Kongresi Bildirileri, 27-29 Ağustos 2007, Isparta, Türkiye, 209.
- Çoruh, S., Çalmaşur, Ö., & Tozlu, G. (2010). Artvin (Yusufeli)'in Tenthredinidae ve Ichneumonidae (Hymenoptera) türleri. Geçmişten Geleceğe Yusufeli Sempozyumu Bildirileri, 9-11 Haziran 2010, Yusufeli, Artvin, Türkiye, 317-321.
- Çoruh, S., Gürbüz, M.F., Kolarov, J., Yurtcan, M., & Boncukçu Özdan, A. (2013). New and Little Known Species of Ichneumonidae (Hymenoptera) for the Turkish Fauna. *Journal of the Entomological Research Society*, 15(3), 71-83.
- Çoruh, S., Kolarov, J., & Çoruh, İ. (2014a). Ichneumonidae (Hymenoptera) from Anatolia. II. *Turkish Journal of Entomology*, 38, 279-290.
- Çoruh, S., Kolarov, J., & Özbeğ, H. (2014b). The fauna of Ichneumonidae (Hymenoptera) of eastern Turkey with zoogeographical remarks and host data. *Journal of Insect Biodiversity*, 2(16), 1-21.
- Çoruh, S., Kolarov, J., & Çoruh, İ. (2018). Ichneumonidae (Hymenoptera) from Anatolia II. *Linzer Biologische Beiträge*, 50(1), 217-224.
- Çoruh, S., Kolarov, J., & Ercelep, Ö. S. (2019). A contribution to the Ichneumoninae (Hymenoptera: Ichneumonidae) of Trabzon, Turkey. *Munis & Entomology Zoology*, 14(2), 584-590.
- Çoruh, S., Kolarov, J., & Ercelep, Ö. S. (2022). A Contribution to the Ichneumonidae (Hymenoptera) of Trabzon, III. Cryptinae. *Atatürk University, Journal of Agricultural Faculty*, 53(1), 8-13.
- Eroğlu, F., Kıraç, A. & Birol, O. (2011). A Faunistic study on Ichneumonidae (Hymenoptera) in Türkmen Mountain, Turkey. *Linzer Biologische Beiträge*, 43(2), 1219-1228.
- Fahringer, J. (1922). Hymenopterologische Ergebnisse einer wissenschaftlichen Studienreise nach der Türkei und Kleinasien (mit Ausschluß des Amanusgebirges). *Archiv für Naturgeschichte*, A(88), 149-222.
- Gallai, N., Salles, J. M., Settele, J., & Vaissiere, B. E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics*, 68, 810-821.
- Gauld, I. D. & Bolton, B. (1988). *The Hymenoptera*. Oxford University Press/British Museum National History. Oxford University Press, Oxford, New York & Toronto. 332 pp.
- Gürbüz, M. F. (2004). *Isparta İli Ichneumonidae (Hymenoptera) Familyası Türleri Üzerine Faunistik ve Sistematis Çalısmalar*. Süleyman Demirel Üniversitesi, Fen Bilimleri Enstitüsü, Isparta, 68 s.
- Gürbüz, M. F. (2005). A survey of the Ichneumonidae (Hymenoptera) of Isparta in Turkey. *Linzer Biologische Beiträge*, 39(2), 1809-1912.
- Gürbüz, M.F., Kirtay, H., & Birol, O. (2009). A study of Ichneumonidae (Hymenoptera) of Kasnak Oak Forest Nature Reserve in Turkey with new records. *Linzer Biologische Beiträge*, 41(2), 1985-2003.
- Gürbüz, M.F., Ljubomirov, T., Kolarov, J., Yurtcan, M., Tabur, M.A., Çoruh, S., & Buncukçu, A. (2008). Investigation of the Ichneumonidae, Ampulicidae, Crabronidae and Sphecidae (Hymenoptera, Insect) Fauna in Natural Protection Zones of East Mediteranean Region in Turkey. *Tübitak TBAGU/ 168 (106T189) No'luproje*, 127 s.
- Jonsson, M., Wratten, S.D., Landis, D.A., & Gurr, G.M. (2008). Recent advances in conservation biological control of arthropods by arthropods. *Biological Control*, 45, 172-175.
- Kirtay, H. (2008). An Investigation on Ichneumonidae (Hymenoptera) Fauna in Kasnak Oak (*Quercus vulcanica* Boiss. and Heldr. ex Kotschy) Forest Nature Protect Area, Isparta. Süleyman Demirel Üniversitesi, Institute of Science, Isparta, Türkiye, 77 s.
- Kıraç, A. & Gürbüz, M. F. (2020). Honaz Dağı Milli Parkı Ichneumonidae (Insecta, Hymenoptera) Faunası. *Bilge International Journal of Science and Technology and Research*, 4(2), 150-159.
- Kolarov, J. & Beyarslan, A. (1994). Investigations on the Ichneumonidae (Hym.) Fauna of Turkey. 1. Pimplinae and Tryphoninae. *Turkish Journal of Entomology*, 18(3), 133-140.

### New Records on the Ichneumonidae Fauna

- Kolarov, J. (1995). A catalogue of the Turkish Ichneumonidae (Hymenoptera). *Zeitschrift für Entomologie*, 7, 137-188.
- Kolarov, J. & Gürbüz, M F. (2004). A study of the Turkish Ichneumonidae (Hymenoptera) I. Pimplinae. *Linzer Biologische Beiträge*, 36(2), 841-845.
- Kolarov, J. & Çalmaşur, Ö. (2011). A study of Ichneumonidae (Hymenoptera) from North Eastern Turkey. *Linzer Biologische Beiträge*, 43(1), 777-782.
- Kolarov, J., Yurtcan, M. & Beyarslan, A. (1997a). New and rare Ichneumonidae (Hym.) from Turkey. 1. Pimplinae, Tryphoninae, Phygadeuontinae, Banchinae and Ctenopelmatinae. *Acta Entomologica Bulgarica*, 3(3/4), 10-12.
- Kolarov, J., Beyarslan, A., & Yurtcan, M. (1997b). Ichneumonidae (Hym.) from the Gökçeada and Bozcaada islands-Turkey. *Acta Entomologica Bulgarica*, 3(3/4), 13-15.
- Kolarov, J., Özbek, H. & Yıldırım, E. (1999). New distributional data of the Turkish Ichneumonidae (Hymenoptera). I. Pimplinae and Tryphoninae. *Journal of the Entomological Research Society*, 1(2), 9-15.
- Kolarov, J., Yurtcan, M., & Beyarslan, A. (2002). Ichneumonidae Species of the Turkish Aegean Region. Parasitic Wasps: Evolution, Systematics. *Biodiversity and Biological Control*, 299- 305.
- Kolarov, J., Çoruh, S., & Çoruh, İ. (2014a). Ichneumonidae (Hymenoptera) from Anatolia. III. *Turkish Journal of Entomology*, 38, 377-388.
- Kolarov, J., Çoruh, S., & Çoruh, İ. (2015). Oxytorinae, a new subfamily for the Turkish fauna (Hymenoptera: Ichneumonidae). *Turkish Journal of Zoology*, 39, 832-835.
- Kolarov, J., Çoruh, S., & Çoruh, İ. (2016). Contribution to the knowledge of the Ichneumonidae (Hymenoptera) fauna of Turkey from northeastern Anatolia, Part I. *Turkish Journal of Zoology*, 40(1), 40-56.
- Kolarov, J., Çoruh, S., Çoruh, İ. (2017). A study of Ichneumonidae (Hymenoptera) from Northeastern Anatolia III, with new records and description male of *Temelucha pseudocaudata* Kolarov, 1982. *Turkish Journal of Entomology*, 41(2): 125-146.
- Kolarov, J., Çoruh, S., & Çoruh, İ. (2020). Ichneumonidae (Hymenoptera) from Anatolia III. *Atatürk University Journal of Agricultural Faculty*, 5(2), 162-168.
- Kolarov, J., Çoruh, S., & Ercelep, Ö. S. (2021). A contribution to the Ichneumonidae (Hymenoptera) of Trabzon, Turkey. II. Campopleginae. *Munis Entomology & Zoology*, 16(2), 745-750.
- Kolarov, J., Çoruh, S., Yurtcan, M., & Gürbüz, M. F. (2009). A study of Metopiinae from Turkey with description of a new species (Hymenoptera: Ichneumonidae). *Zoology in the Middle East*, 46, 75-82.
- Kolarov, K., Yıldırım, E., Çoruh, S., & Yüksel, M. (2014b). Contribution to the knowledge of the Ichneumonidae (Hymenoptera) fauna of Turkey, *Zoology in the Middle East*, 60(2), 154-161.
- Kolarov, J., Pekel, S., Özbek, H., Yıldırım, E., & Çalmaşur, Ö. (2000). New distributional data of Turkish Ichneumonidae (Hymenoptera). III. The subfamily Ophioninae. *Türkiye IV Entomoloji Kongresi*, 12-15 Eylül 2000, Kuşadası-Aydın, Türkiye, 349-356.
- Narmanlioğlu, H.K. & Coruh, S. (2017). Parasitoids of the apple ermine moth, *Yponomeuta malinellus* Zeller, 1838 (Lepidoptera: Yponomeutidae), in the Çoruh Valley, Erzurum Province, Turkey. *Turkish Journal of Entomology*, 41(4), 357-365.
- Okyar, Z. & Yurtcan, M. (2007). Phytophagous Noctuidae (Lepidoptera) of the Western Black Sea Region and their ichneumonid parasitoids. *Entomofauna*, 28, 377-388.
- Öncüler, C. (1991). *Türkiye Bitki Zararlısı Böceklerinin Parazit ve Predatör Kataloğu*. Ege Üniversitesi, Ziraat Fakültesi Yayınları, 505, 354.
- Özdemir, Y. & Kilinçer, H. (1990). The species of Pimplinae and Ophioninae from Central Anatolia. Proceedngs of the Second Turkish National Congress of Biological Control, 1990, Ankara, Turkey, 309-318.
- Özbek, H. & Çoruh, S. (2012). Larval parasitoids and larval diseases of *Malacosoma neustria* L. (Lepidoptera: Lasiocampidae) detected in Erzurum Province, Turkey. *Turkish Journal of Zoology*, 36(4), 447-459.

- Özdan, A. (2014). *Gelincik Dağı Tabiat Parkı ve Kovada Gölü Milli Parkı (İsparta) Ichneumonidae (Hymenoptera) Faunası*. Süleyman Demirel Üniversitesi, Fen Bilimleri Enstitüsü, Isparta, 150 s.
- Özdan, A., Gürbüz, M.F. (2019). Ichneumonidae (Hymenoptera) fauna of Kovada Lake National Park, Isparta, Turkey. *Turkish Journal of Entomology*, 43(3): 301-312.
- Rasplus, J-Y., Villemant, C., Paiva, M.R., Delvare, G., Roques, A. (2010). Hymenoptera. Chapter 12. In Roques A et al (Eds.). *Arthropod invasions in Europe* (page range). *BioRisk*, 4(2), 669-776.
- Riedel, M., Diller E., & Çoruh, S. (2018). New Contributions to the Ichneumoninae (Hymenoptera, Ichneumonidae) from Turkey. *Journal of the Entomological Research Society*, 20(1): 57-70.
- Riedel, M., Kolarov, J., Çoruh, S. & Özbek, H. (2014). A contribution to the Mesochorinae (Hymenoptera: Ichneumonidae) of Turkey. *Zoology in the Middle East*, 60, 217-221.
- Sarı, Ü. & Çoruh, S. (2018). Ichneumonidae (Hymenoptera) from Northeastern Anatolia Region (Erzurum, Aşkale). *Turkish Journal of Entomology*, 42(3): 215-228.
- Schwarz, M. (2020). Zer Kenntnis der palaarktischen *Meringopus* -Arten (hymenoptera, Ichneumonidae, Cryptinae). *Linzer Biologische Beiträge*, 52(1): 583-682.
- Teymuroğlu, E. & Çoruh, S. (2021). Ichneumonidae (Hymenoptera) species associated with sugar beet plants occurring in Erzincan (Çayırlı) of Turkey. *Munis Entomology & Zoology*, 16(2), 985-990.
- Tscharntke, T., Bommarco, R., Clough, Y., Crist, T.O., & Kleijn, D. (2007). Conservation biological control and enemy diversity on a landscape scale. *Biological Control*, 43, 294-309.
- Wahl, D.B. & Micheal, J.S. (1993). Family Ichneumonidae. In Goulet H., Huber J. (Eds.), *The Hymenoptera of the world, an identification guide to families* (pp. 395-442) , Canada Agriculture, Canada.
- Vas, Z. (2019a). Contributions to the taxonomy, identification, and biogeography of the Western Palaearctic species of *Campoletis* Förster (Ichneumonidae: Campopleginae). *Zootaxa*, 4565(3), 373-382.
- Vas, Z. (2019b). New species and new records of Ichneumon wasps from the Eastern Mediterranean and the Black Sea Regions (Hymenoptera: Ichneumonidae). *Acta Zoologica Academiae Scientiarum Hungaricae*, 65(1), 19-30.
- Yaman, G. (2014). Türkiye Tryphoninae (Hymenoptera: Ichneumonidae) türlerinin kontrol listesi. Trakya University, Fen Bilimleri Enstitüsü, Edirne, 88 s.
- Yu, D. S. Ki, Achterberg, C. Van, & Horstmann, K. (2016). Taxapad 2016, Ichneumonoidea 2015. Database on flash-drive. [www.taxapad.com](http://www.taxapad.com), Nepean, Ontario, Canada.
- Yurtcan, M. (2004). Trakya Bölgesi Pimplinae (Hymenoptera: Ichneumonidae) Faunasının Taksonomik ve Faunistik Yönden Araştırılması, Trakya Üniversitesi Fen Bilimleri Enstitüsü, Edirne,
- Yurtcan, M. (2007). Ephialtini tribe (Hymenoptera, Ichneumonidae, Pimplinae) of Turkish Thrace region, Entomofauna, 28, 389-404.
- Yurtcan, M. & Beyarslan, A. (2002). The species of Tryphoninae (Hymenoptera: Ichneumonidae) in Turkish Thrace. *Turkish Journal of Zoology*, 26(1), 77-95.
- Yurtcan, M. & Beyarslan, B. (2005). Polysphinctini and Pimplini (Hymenoptera: Ichneumonidae: Pimplinae) from the thrace Region of Turkey. *Fragmenta Faunistica*, 48(1), 63-72.
- Yurtcan, M., Çoruh, S., Kolarov, J., Özdan, A.B., Gürbüz, M.F. & Erkaya, İ. (2021). Ichneumonidae (Hymenoptera) fauna of natural protection areas in the east mediterranean region of Turkey, Part II. *Entomological News*, 129(5), 453-472.